

The $K_L \rightarrow \pi^0 \nu \bar{\nu}$ Decay Beyond the Standard Model

Yuval Grossman

*Stanford Linear Accelerator Center
Stanford University, Stanford, CA 94309***Abstract**

The $K_L \rightarrow \pi^0 \nu \bar{\nu}$ decay is analyzed in a model independent way. When lepton flavor is conserved, this decay mode is a manifestation of CP violating interference between mixing and decay. Consequently, a theoretically clean relation between the measured rate and electroweak parameters holds in any given model.

$K_L \rightarrow \pi^0 \nu \bar{\nu}$ is unique among K decays in several aspects: (a) It is theoretically very clean; (b) it is purely CP violating^{1,2}; and (c) it can be measured in the near future³ even if the rate is as small as the Standard Model prediction. In the Standard Model a measurement of $\Gamma(K_L \rightarrow \pi^0 \nu \bar{\nu})$ provides a clean determination of the Wolfenstein CP violating parameter η or, equivalently, of the Jarlskog measure of CP violation J and, together with a measurement of $\Gamma(K^+ \rightarrow \pi^+ \nu \bar{\nu})$, of the angle β of the unitarity triangle².

Here we explain what can be learned from the $K \rightarrow \pi \nu \bar{\nu}$ decay in a model independent way⁴. We define

$$\lambda \equiv \frac{q}{p} \frac{\bar{A}}{A}, \quad (1)$$

where p and q are the components of interaction eigenstates in mass eigenstates, $|K_{L,S}\rangle = p|K^0\rangle \mp q|\bar{K}^0\rangle$, and $A(\bar{A})$ is the $K^0(\bar{K}^0) \rightarrow \pi^0 \nu \bar{\nu}$ decay amplitude. Then, the ratio between the K_L and K_S decay rates is⁴

$$\frac{\Gamma(K_L \rightarrow \pi^0 \nu \bar{\nu})}{\Gamma(K_S \rightarrow \pi^0 \nu \bar{\nu})} = \frac{1 + |\lambda|^2 - 2\text{Re}\lambda}{1 + |\lambda|^2 + 2\text{Re}\lambda}. \quad (2)$$

In general, a three body final state does not have a definite CP parity. However, if the light neutrinos are purely left-handed, and if lepton flavor is conserved, the final state is CP even (to an excellent approximation)⁴. If lepton flavor is violated, the final state in $K_L \rightarrow \pi^0 \nu \bar{\nu}$ is not necessarily a CP eigenstate; specifically, $K_L \rightarrow \pi^0 \nu_i \bar{\nu}_j$ with $i \neq j$ is allowed. Here, we concentrate on the

Talk given at the first symposium of FCNC, February 19-21, 1997, Santa Monica, CA.

case where the above two conditions are satisfied, so that the final state is purely CP even.

The contributions to the $K_L \rightarrow \pi^0 \nu \bar{\nu}$ decay from CP violation in mixing ($|q/p| \neq 1$) and from CP violation in decay ($|\bar{A}/A| \neq 1$) are negligibly small. The deviation of $|q/p|$ from unity is experimentally measured (by the CP asymmetry in $K_L \rightarrow \pi \ell \nu$) and is $\mathcal{O}(10^{-3})$. The deviation of $|\bar{A}/A|$ from unity is expected to be even smaller⁴. Therefore, $|\lambda| = 1 + \mathcal{O}(10^{-3})$, and the leading CP violating effect is $\text{Im}\lambda \neq 0$, namely interference between mixing and decay. This puts the ratio of decay rates (2) in the same class as CP asymmetries in various B decays to final CP eigenstates, e.g. $B \rightarrow \psi K_S$, where a very clean theoretical analysis is possible⁵.

As a result of this cleanliness, the CP violating phase can be extracted almost without any hadronic uncertainty, even if this phase comes from New Physics. Defining θ to be the relative phase between the $K - \bar{K}$ mixing amplitude and the $s \rightarrow d \nu \bar{\nu}$ decay amplitude, namely $\lambda = e^{2i\theta}$, we get from eq. (2)

$$\frac{\Gamma(K_L \rightarrow \pi^0 \nu \bar{\nu})}{\Gamma(K_S \rightarrow \pi^0 \nu \bar{\nu})} = \frac{1 - \cos 2\theta}{1 + \cos 2\theta} = \tan^2 \theta. \quad (3)$$

In reality, however, it will be impossible to measure $\Gamma(K_S \rightarrow \pi^0 \nu \bar{\nu})$. We can use the isospin relation, $A(K^0 \rightarrow \pi^0 \nu \bar{\nu})/A(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = 1/\sqrt{2}$, to replace the denominator by the charged kaon decay mode:

$$a_{CP} \equiv r_{\text{is}} \frac{\Gamma(K_L \rightarrow \pi^0 \nu \bar{\nu})}{\Gamma(K^+ \rightarrow \pi^+ \nu \bar{\nu})} = \frac{1 - \cos 2\theta}{2} = \sin^2 \theta, \quad (4)$$

where $r_{\text{is}} = 0.954$ is the isospin breaking factor⁶. The ratio (4) may be experimentally measurable as the relevant branching ratios are $\mathcal{O}(10^{-10})$ in the Standard Model² and even larger in some of its extensions.

Eq. (4) implies that a measurement of a_{CP} will allow us to determine the CP violating phase θ without any information about the magnitude of the decay amplitudes. Also, using $\sin^2 \theta \leq 1$ and $\tau_{K_L}/\tau_{K^+} = 4.17$, we get the model independent bound

$$\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) < 1.1 \times 10^{-8} \left(\frac{\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})}{2.4 \times 10^{-9}} \right). \quad (5)$$

This bound is much stronger than the direct experimental upper bound⁷ $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) < 5.8 \times 10^{-5}$.

New Physics can modify both the mixing and the decay amplitudes. $\varepsilon = \mathcal{O}(10^{-3})$ implies that any new contribution to the mixing amplitude carries almost the same phase as the Standard Model one. On the other hand, the

upper bound⁸ $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) < 2.4 \times 10^{-9}$, which is much larger than the Standard Model prediction², allows New Physics to dominate the decay amplitude (with an arbitrary phase). We conclude that a significant modification of a_{CP} can only come from New Physics in the decay amplitude. For example, in models with extra quarks, the decay amplitudes can be dominated by tree level Z -mediated diagrams⁴.

In superweak models, all CP violating effects appear in the mixing amplitudes. Then, CP violation in $K_L \rightarrow \pi^0 \nu \bar{\nu}$ should be similar in magnitude to that in $K_L \rightarrow \pi \pi$. In models of approximate CP symmetry, all CP violating effects are small. Both scenarios predict then $a_{CP} = \mathcal{O}(10^{-3})$, in contrast to the Standard Model prediction, $a_{CP} = \mathcal{O}(1)$. In other words, a measurement of $a_{CP} \gg 10^{-3}$ (and, in particular, $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) \gtrsim \mathcal{O}(10^{-11})$) will exclude these two scenarios of New Physics in CP violation.

In the Standard Model there are two clean ways to determine the unitarity triangle: (1) CP asymmetries in B^0 decays⁵; and (2) the combination of $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ ². In general, New Physics will affect both determinations. Moreover, it is very unlikely that the modification of the two methods will be the same. Consequently, a comparison between these two clean determinations will be a very powerful tool to probe CP violation beyond the Standard Model. Because of the very small theoretical uncertainties in both methods even a small new physics effect can be detected. In practice, we will be limited only by the experimental sensitivity.

In conclusion: a measurement of $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ is guaranteed to provide us with valuable information. It will either give a new clean measurement of CP violation or indicate lepton flavor violation.

Acknowledgments. I thank Yossi Nir for collaboration on this work. Y.G. is supported by the Department of Energy under contract DE-AC03-76SF00515.

1. L.S. Littenberg, Phys. Rev. **D 39** (1989) 3322.
2. G. Buchalla and A.J. Buras, Nucl. Phys. **B 400** (1993) 225; Phys. Rev. **D 54** (1996) 6782; A.J. Buras, Phys. Lett. **B 333** (1994) 476; G. Buchalla, these proceedings.
3. T. Inagaki, these proceedings; D. Bryman, these proceedings; K. Arisaka, these proceedings.
4. Y. Grossman, and Y. Nir, hep-ph/9701313, to appear in Phys. Lett. **B**.
5. See, e.g. Y. Nir and H.R. Quinn, Ann. Rev. Nucl. Part. Sci. **42** (1992) 211.
6. W. Marciano and Z. Parsa, Phys. Rev. **D 53** (1996) 1.
7. M. Weaver *et al.*, E799 Collaboration, Phys. Rev. Lett. **72** (1994) 3758.
8. S. Adler *et al.*, BNL 787 Collaboration, Phys. Rev. Lett. **76** (1996) 1421.